

Drugs in interaction	Resulting side effects
Quinolones + macrolides	Prolongation of QTc, arrhythmias
ACE inhibitors + spironolactone	Hyperkalemia
SSRIs + triptans	Serotonin syndrome
NSAIDs + glucocorticoids	Gastrointestinal bleeding
Gentamicin + ampicillin	Increased bactericidal effect
Acetaminophen + aspirin	Increased pain control
Sildenafil + nitroglycerine	Severe hypotension
MAO inhibitors + high tyramine foods (cheese, wine, cured meats, bananas, etc.)	Potentially fatal increase in blood pressure, headache
Warfarin + Vit K-rich foods (kale, broccoli, etc.)	Reduced effect of warfarin causing blood clots

Drugs in interaction	Common uses	Resulting side effects
Ginkgo biloba	Warfarin, clopidogrel, aspirin	Risk of bleeding
Licorice	Warfarin Digoxin	Risk of clotting, Digoxin toxicity
St.John's wort	Warfarin Multiple interactions with cardiac drugs, statins, immunosuppressants, antidepressants, benzodiazepines, antiepileptics, birth control pills, anti-HIV drugs	Risk of clotting Decreased effect, increased toxicity

Drugs in interaction	Resulting side effects
ACE inhibitors + potassium supplements	Hyperkalemia
Digoxin + amiodarone	Digoxin toxicity
Digoxin + verapamil	Bradycardia, heart block, digoxin toxicity
Theophylline + quinolones	Theophylline toxicity, seizures
Warfarin + macrolides/quinolones/sulfa antibiotics	Bleeding
Warfarin + NSAIDs	Gastrointestinal bleeding
Warfarin + phenytoin	Warfarin toxicity, phenytoin toxicity
Simvastatin + fluconazole	Increased simvastatin levels, liver toxicity
Levothyroxine + omeprazole	Decreased absorption of levothyroxine
Metronidazole + alcohol	Nausea, vomiting, flushing, headache